

B_s Mixing and New Physics in hadronic $b \rightarrow s\bar{q}q$ transitions

Alakabha Datta ^{a1}

*a: Department of Physics, University of Toronto,
60 St. George Street, Toronto, ON, Canada M5S 1A7*

(February 2, 2008)

Abstract

We study the implication of the recent ΔM_s measurement on $b \rightarrow s\bar{q}q$ transitions. We show that it is possible, in the presence of a flavour symmetry, that the phase in B_s mixing may be unobservable even with new CP odd phases in $b \rightarrow s$ transitions. These phases may then produce new CP odd effects in certain $b \rightarrow s\bar{q}q$ transitions like $B \rightarrow K\pi$ but not in others like $B_d \rightarrow \phi K_s$. Working in a two higgs doublet model of new physics we discuss the allowed NP contribution to $B \rightarrow K\pi$ and $B_d \rightarrow \phi K_s$ decays with the new ΔM_s measurement.

¹datta@physics.utoronto.ca

1 Introduction

The $B_s^0 - \bar{B}_s^0$ mass difference, recently measured by the DØ [1] and CDF [2] collaborations, is given by,

$$\begin{aligned} 17 \text{ ps}^{-1} < \Delta M_s < 21 \text{ ps}^{-1} \quad (90\% \text{ CL, DØ}), \\ \Delta M_s = (17.33_{-0.21}^{+0.42} \pm 0.07) \text{ ps}^{-1} \quad (\text{CDF}) \end{aligned} \quad (1)$$

This result is consistent with the SM predictions, which are estimated as $21.3 \pm 2.6 \text{ ps}^{-1}$ by the UTfit group [3] and $20.9_{-4.2}^{+4.5} \text{ ps}^{-1}$ $21.7_{-4.2}^{+5.9} \text{ ps}^{-1}$ by the CKMfitter group [4]. The implication of this measurement have been studied in several recent papers [5, 6, 7]. Given the hadronic uncertainties in the SM prediction, along with additional uncertainties when NP is included, the present measurement of ΔM_s does not provide a strong constraint on NP[6, 7].

A measurement of a large phase in B_s mixing will be a definite signal of NP as the SM prediction for this phase is $1.045_{-0.057}^{+0.061}$ degrees [8] and is tiny. NP phases should then also appear in $\Delta B = 1$, $b \rightarrow s\bar{q}q$ ($q = u, d, s$) hadronic transitions. However a measurement of no significant NP phase in B_s mixing will not necessarily rule out new NP weak phases in $b \rightarrow s\bar{q}q$ transitions. In fact Ref. [9] finds that combining the measurements of the mass difference ΔM_s and that of the semileptonic CP asymmetry one can constrain the B_s mixing to be small. In this paper we show that one can construct a flavour symmetry that causes new CP odd phase, from the $b \rightarrow s$ vertex, in B_s mixing to be unobservable but allows new CP odd effects in $\Delta B = 1$, $b \rightarrow s\bar{q}q$ transitions.

There is now a lot of data on rare FCNC hadronic decays from various experiments. The data, specially involving $b \rightarrow s\bar{q}q$ transitions, have been somewhat difficult to understand within the standard model(SM). First, within the SM, the measurement of the CP phase $\sin 2\beta$ in $B_d^0(t) \rightarrow J/\psi K_s$ should be approximately equal to that in decays dominated by the quark-level penguin transition $b \rightarrow s\bar{q}q$ ($q = u, d, s$) like $B_d^0(t) \rightarrow \phi K_s$, $B_d^0(t) \rightarrow \eta' K_s$, $B_d^0(t) \rightarrow \pi^0 K^0$, etc. However, there is a difference between the measurements of $\sin 2\beta$ in the $b \rightarrow s$ penguin dominated modes ($\sin 2\beta = 0.50 \pm 0.06$) and that in $B_d^0(t) \rightarrow J/\psi K_s$ ($\sin 2\beta = 0.685 \pm 0.032$) [10, 11, 12]. Note that the $\sin 2\beta$ number for the $b \rightarrow s$ penguin dominated modes is the average of several modes. The effect of new physics can be different for different final hadronic states and so the individual $\sin 2\beta$ measurements for the different modes are important. Second, the latest data on $B \rightarrow \pi K$ decays (branching ratios and various CP asymmetries) appear to be inconsistent with the SM [13, 14].² Third, within the SM, one expects no triple-product asymmetries in $B \rightarrow \phi K^*$ [16], but BaBar has measured such an effect at 1.7σ level [17]. There are also polarization anomalies where the decays $\bar{B}_d^0 \rightarrow \phi K^*$ and $B^- \rightarrow \rho^- K^*$ appear to

²A cleaner test of the SM could be provided by looking at the quasi-exclusive decays $B \rightarrow KX$ [15] rather than the exclusive $B \rightarrow K\pi$ decays.

have large transverse polarization amplitudes in conflict with naive SM expectations [12, 18, 19, 20, 21]. While these deviations certainly do not unambiguously signal new physics(NP), they give reason to speculate about NP explanations of the experimental data. Hence, it is interesting to study the implication of the measurement of ΔM_s for $b \rightarrow s\bar{q}q$ transitions. For instance, one could check if the NP fits to the $B \rightarrow K\pi$ decays presented in Ref. [14] are consistent with ΔM_s measurements.

The paper is organized in the following manner: In the next section we study NP phases in B_s mixing and a flavour symmetry under which the new phase in B_s mixing remains hidden even with NP with non trivial weak phases. We then discuss how these new weak phases affect hadronic $b \rightarrow s\bar{q}q$ transitions. In the following section we present details of a two higgs model already presented in Ref. [22] to demonstrate explicitly some of the general ideas of sec.2 and to quantitatively study the implication of ΔM_s measurements for $B \rightarrow K\pi$ and $B_d^0 \rightarrow \phi K_s$ decays. Finally in sec.4 we present our conclusions.

2 Phase of B_s Mixing

The mass difference ΔM_s^{SM} is calculated in the standard model from the $B_s^0 - \overline{B}_s^0$ box diagram, dominated by t -quark exchange:

$$\begin{aligned} M_{12}^s &= \frac{G_F^2}{6\pi^2} M_W^2 M_{B_s} \left(f_{B_s}^2 \hat{B}_{B_s} \right) \hat{\eta}_{B_s} y_t f_2(y_t) |V_{ts}^* V_{tb}|^2, \\ \Delta M_s^{SM} &= |M_{12}^s|, \end{aligned} \quad (2)$$

where $y_t = m_t^2/M_W^2$, and the function f_2 is the Inami-Lim function [23]. $\hat{\eta}_{B_s}$ is the QCD correction and $f_{B_s}^2 \hat{B}_{B_s}$ represents the hadronization of the B_s mixing operator in the SM.

New physics can generate operators of the type

$$H_{NP}^{\Delta B=2} \sim \sum \frac{c_{ij}}{\Lambda^2} (\bar{s}\Gamma_i b)(\bar{s}\Gamma_j b), \quad (3)$$

that will contribute to ΔM_s . Here Λ is the scale of new physics and c_{ij} are the strength of the operators. If the scale of NP is around a TeV the size c_{ij} must be $\sim |V_{ts}|^2$ to produce effects that can be of the same size as the experimental measurements [5, 6, 7]. Experimental information on ΔM_s can constrain the c_{ij} but this process is complicated by the fact that there can be more than one operator in Eq. 3 and the fact that the different NP operators can hadronize differently than the SM operator. Assuming for the sake of simplification that only two operators contribute in Eq. 3, we can write

$$\begin{aligned} M_{12}^s &= M_{12}^{s,SM} + M_{12}^{s,NP}, \\ M_{12}^{s,NP} &= \frac{\langle B_s | H_{NP}^{\Delta B=2} | \overline{B}_s^0 \rangle}{M_{B_s}}, \end{aligned}$$

$$\Delta M_s = \Delta M_s^{SM} \left| 1 + r_1 \eta_1 e^{i\phi_1} + r_2 \eta_2 e^{i\phi_2} \right|, \quad (4)$$

where r_1 and r_2 represent the different hadronization of the NP operators relative to the SM operators. The quantities $\eta_{1,2}$ may be related to c_{ij} in Eq. 3 in a specific model of NP. One may combine to write the NP amplitudes as a single complex number as

$$\begin{aligned} \Delta M_s &= \Delta M_s^{SM} \left| 1 + \eta e^{i\phi} \right|, \\ \eta e^{i\phi} &= r_1 \eta_1 e^{i\phi_1} + r_2 \eta_2 e^{i\phi_2}, \end{aligned} \quad (5)$$

where the relation between η and the size of the NP operators is complicated by SM uncertainties, QCD effects of NP operators and more than one NP amplitude. As indicated in the introduction the measurement of the phase in B_s mixing is a good place to look for NP. From the general structure of Eq. 5 it is trivially true that if $\phi_{1,2} = 0$ then there is no new NP weak phase in B_s mixing. However, even with $\phi_{1,2} \neq 0$ it is possible that the effective phase ϕ in Eq. 5 vanishes. This may be due to accidental cancellation or flavour symmetries where the NP contributions may conspire to produce a small or no effective phase ϕ even though ϕ_1 and ϕ_2 may be large.

In this section we study a flavour symmetry under which the new phase in B_s mixing remain hidden even with NP with nontrivial weak phases. We then study the implications for such a scenario for $b \rightarrow s\bar{q}q$ transitions.

The condition that the effective phase $\phi = 0, \pi$ in Eq. 5 is

$$\begin{aligned} M_{12}^{s,NP} &= M_{12}^{s,NP*} \Rightarrow \\ \langle B_s | H_{NP}^{\Delta B=2} | \overline{B}_s^0 \rangle &= \langle \overline{B}_s^0 | (H_{NP}^{\Delta B=2})^\dagger | B_s \rangle. \end{aligned} \quad (6)$$

Since $CP|B_s\rangle \sim |\overline{B}_s^0\rangle$ the condition in Eq. 6 is satisfied with the requirement

$$(H_{NP}^{\Delta B=2})_{CP} = (H_{NP}^{\Delta B=2})^\dagger, \quad (7)$$

where $(H_{NP}^{\Delta B=2})_{CP}$ is just the CP transformed $H_{NP}^{\Delta B=2}$. However it is easy to see, and is well known, that the requirement in Eq. 7 just leads to the trivial possibility that all weak phases $\phi_{1,2}$ in Eq. 4(Eq. 5) vanish. We will demonstrate this with two models of NP that we call vector-axial vector model (V/A) and the scalar-pseudoscalar model(S/P). We write the general NP lagrangian as

$$\begin{aligned} H_{NP,A}^{\Delta B=2} &= C_{LL,A} O_{LL,A} + C_{LR,A} O_{LR,A} + C_{RL,A} O_{RL,A} + C_{RR,A} O_{RR,A}, \\ O_{LL,A} &= \bar{s}\gamma_A(1 - \gamma_5)b\bar{s}\gamma^A(1 - \gamma_5)b, \\ O_{LR,A} &= \bar{s}\gamma_A(1 - \gamma_5)b\bar{s}\gamma^A(1 + \gamma_5)b, \\ O_{RL,A} &= \bar{s}\gamma_A(1 + \gamma_5)b\bar{s}\gamma^A(1 - \gamma_5)b, \\ O_{RR,A} &= \bar{s}\gamma_A(1 + \gamma_5)b\bar{s}\gamma^A(1 + \gamma_5)b, \end{aligned} \quad (8)$$

where $A = S, V$ and $\gamma^A = 1, \gamma^\mu$ for the S/P and V/A NP models. Now one easily checks that $(O_{X,Y})_{CP} = O_{X,Y}^\dagger$ where $X, Y = L, R$ for the operators in Eq. 8 for both V/A and S/P models of NP. The condition in Eq. 7 then leads to the fact that the coefficients $C_{X,Y}$ in Eq. 8 are real and we get the trivial scenario that all NP weak phases in $H_{NP}^{\Delta B=2}$ vanish.

However we show that the condition in Eq. 6 can also be satisfied with a flavour symmetry. Let us now define a flavour transformation, T , such that

$$\begin{aligned} T|s\rangle &= |b\rangle, \\ T|b\rangle &= |s\rangle. \end{aligned} \quad (9)$$

This is nothing but the $s-b$ interchange transformation discussed in Ref [22] which is the quark equivalent of the $\mu-\tau$ interchange symmetry used in the study of neutrino mixing [24, 25]. From Eq. 9 we obtain

$$\begin{aligned} T|B_s\rangle &= \xi|\overline{B}_s^0\rangle, \\ T|\overline{B}_s^0\rangle &= \xi|B_s\rangle, \\ TH_{NP}^{\Delta B=2}T^\dagger &= T\left[\sum\frac{c_{ij}}{\Lambda^2}(\bar{s}\Gamma_ib)(\bar{s}\Gamma_jb)\right]T^\dagger, \\ &= \sum\frac{c_{ij}}{\Lambda^2}(\bar{b}\Gamma_is)(\bar{b}\Gamma_js), \end{aligned} \quad (10)$$

where $|\xi|^2 = 1$.

Hence to recover the condition in Eq. 6 we demand that invariance of $H_{NP}^{\Delta B=2}$ under the $s-b$ flavour symmetry implies

$$\begin{aligned} TH_{NP}^{\Delta B=2}T^\dagger &= (H_{NP}^{\Delta B=2})^\dagger \Rightarrow \\ \sum\frac{c_{ij}}{\Lambda^2}(\bar{b}\Gamma_is)(\bar{b}\Gamma_js) &= \sum_i\frac{c_{ij}^*}{\Lambda^2}(\bar{b}\gamma_0\Gamma_j^\dagger\gamma_0s)(\bar{b}\gamma_0\Gamma_i^\dagger\gamma_0s). \end{aligned} \quad (11)$$

Now writing $\gamma_0\Gamma_{ij}^\dagger\gamma_0 = \pm\Gamma_{i'j'}$ the relation in Eq. 11 then translates, in general, to

$$c_{ij}^* = \pm c_{i'j'} \quad (12)$$

For the particular case of the V/A model of NP $\Gamma_{ij} = \gamma_\mu(1 \pm \gamma_5)$ and we have $\gamma_0\Gamma_{ij}^\dagger\gamma_0 = \Gamma_{ij}$ leading to $c_{ij}^* = c_{ij}$ and so the coefficient of each individual operator in $L_{NP,V}$ is real. This is easy to understand as in this case invariance under $s-b$ flavour symmetry is the same as requiring invariance under CP. It is possible to impose a weaker requirement for $s-b$ symmetry as

$$\langle \overline{B}_s^0 | TH_{NP}^{\Delta B=2}T^\dagger | B_s \rangle = \langle \overline{B}_s^0 | (H_{NP}^{\Delta B=2})^\dagger | B_s \rangle. \quad (13)$$

This then leads to, using Eq. 8,

$$\begin{aligned} &\langle \overline{B}_s^0 | C_{LL,V}O_{LL,V}^\dagger + C_{LR}O_{LR,V}^\dagger + C_{RL,V}O_{RL}^\dagger + C_{RR,V}O_{RR,V}^\dagger | B_s \rangle = \\ &\langle \overline{B}_s^0 | C_{LL,V}^*O_{LL,V}^\dagger + C_{LR}^*O_{LR,V}^\dagger + C_{RL,V}^*O_{RL}^\dagger + C_{RR,V}^*O_{RR,V}^\dagger | B_s \rangle. \end{aligned} \quad (14)$$

Now as $\langle \overline{B}_s^0 | O_{LL,V}^\dagger | B_s \rangle = \langle \overline{B}_s^0 | O_{RR,V}^\dagger | B_s \rangle$ and $\langle \overline{B}_s^0 | O_{LR,V}^\dagger | B_s \rangle = \langle \overline{B}_s^0 | O_{RL,V}^\dagger | B_s \rangle$ using parity transformation, we see that Eq. 13 is also satisfied with the non trivial solution

$$\begin{aligned} C_{LL,V}^* &= C_{RR,V}, \\ C_{LR,V}^* &= C_{RL,V}. \end{aligned} \quad (15)$$

For the S/P models $s - b$ symmetry of the lagrangian is possible with complex coefficients in the lagrangian and here the relation in Eq. 12 can be written as,

$$\begin{aligned} C_{LL,S}^* &= C_{RR,S}, \\ C_{LR,S}^* &= C_{RL,S}. \end{aligned} \quad (16)$$

Hence $H_{NP,A}^{\Delta B=2}$ has the form, using Eq. 15 and Eq. 16 in Eq. 8,

$$H_{NP,A}^{\Delta B=2} = |a| \left[e^{-2i\phi} O_{LL,A} + e^{2i\phi} O_{RR,A} \right] + |b| \left[e^{-2i\psi} O_{LR,A} + e^{2i\psi} O_{RL,A} \right]. \quad (17)$$

We now ask the question if the phases ϕ and ψ in Eq. 17 can produce CP odd effects in $b \rightarrow s\bar{q}q$ transitions. To answer this question we consider models of NP where the new interaction arises at tree level from the exchange of a heavy field.

Assuming that $H_{NP,A}^{\Delta B=2}$ in Eq. 8 arises after integrating a heavy field then we can write

$$\begin{aligned} H_{NP,A}^{\Delta B=2} &\sim J_{sb}^2, \\ J_{sb,A} &= [\alpha \bar{s} \gamma_A (1 - \gamma_5) b + \beta \bar{s} \gamma_A (1 + \gamma_5) b], \end{aligned} \quad (18)$$

where $A = S, V$ and $\gamma^A = 1, \gamma^\mu$ for the S/P and V/A NP models. The requirements from Eq. 15 and Eq. 16 then allow us to write

$$J_{sb,A} = |\alpha| \left[e^{i\phi} \bar{s} \gamma_A (1 + \gamma_5) b \pm e^{-i\phi} \bar{s} \gamma_A (1 - \gamma_5) b \right]. \quad (19)$$

The relative sign between the terms in Eq. 19 would correspond to a scalar or a pseudoscalar exchange in the S/P model of new physics. We will study such a model in details in the next section.

We can now write Eq. 18 as

$$H_{NP,A}^{\Delta B=2} = |\alpha|^2 \left[e^{-2i\phi} O_{LL,A} \pm O_{LR,A} \pm O_{RL,A} + e^{2i\phi} O_{RR,A} \right]. \quad (20)$$

Even though the phase ϕ in $H_{NP,A}^{\Delta B=2}$ is not observable in B_s mixing its effect may be observed in $\Delta B = 1, b \rightarrow s\bar{q}q$ transitions. We can write the effective Hamiltonian that generates the $b \rightarrow s\bar{q}q$ transition as

$$\begin{aligned} H_{eff}^{\Delta B=1} &\sim J_{sb} J_{qq}, \\ J_{sb,A} &= |\alpha| \left[e^{i\phi} \bar{s} \gamma_A (1 + \gamma_5) b \pm e^{-i\phi} \bar{s} \gamma_A (1 - \gamma_5) b \right], \\ J_{qq,A} &= \left[c_R \bar{q} \gamma^A (1 + \gamma_5) q \pm c_L^* \bar{q} \gamma^A (1 - \gamma_5) q \right]. \end{aligned} \quad (21)$$

Here $c_{L,R}$ are in general complex. We will first consider the case $c_L = c_R = c$ as this relation is satisfied in the model to be considered in the next section. Writing $c = |c|e^{i\phi_c}$ we can rewrite H_{eff} as

$$\begin{aligned}
H_{eff}^{\Delta B=1} &\sim |\alpha||c| \left[e^{i\phi_+} H_{RR,A} + e^{-i\phi_+} H_{LL,A} \pm e^{i\phi_-} H_{RL} \pm e^{-i\phi_-} H_{LR} \right], \\
H_{LL,A} &= \bar{s}\gamma_A(1 - \gamma_5)b\bar{q}\gamma^A(1 - \gamma_5)q, \\
H_{RL,A} &= \bar{s}\gamma_A(1 + \gamma_5)b\bar{q}\gamma^A(1 - \gamma_5)q, \\
H_{LR,A} &= \bar{s}\gamma_A(1 - \gamma_5)b\bar{q}\gamma^A(1 + \gamma_5)q, \\
H_{RR,A} &= \bar{s}\gamma_A(1 + \gamma_5)b\bar{q}\gamma^A(1 + \gamma_5)q,
\end{aligned} \tag{22}$$

where $\phi_{\pm} = \phi \pm \phi_c$.

To demonstrate our point we can use factorization to calculate nonleptonic decays. Let us consider the decay $B^- \rightarrow K^- \pi^0$ and using the currents associated with the relative negative sign in $J_{sb,qq}$, we can write

$$\begin{aligned}
\langle K^- \pi^0 | H_{eff} | B^- \rangle &\sim |\alpha||c| \langle K^- | e^{i\phi} \bar{s}\gamma_A(1 + \gamma_5)b - e^{-i\phi} \bar{s}\gamma_A(1 - \gamma_5)b | B^- \rangle \\
&\quad \langle \pi^0 | e^{i\phi_c} \bar{d}\gamma^A(1 + \gamma_5)d - e^{-i\phi_c} \bar{d}\gamma^A(1 - \gamma_5)d | 0 \rangle, \\
&\sim 4|\alpha||c|e^{i\pi/2} \sin \phi \cos \phi_c \langle K^- | \bar{s}\gamma_A b | B^- \rangle \\
&\quad \langle \pi^0 | \bar{d}\gamma^A \gamma_5 d | 0 \rangle.
\end{aligned} \tag{23}$$

We observe that an effective weak phase of $\pi/2$ appears in the $B \rightarrow K\pi$ amplitude but this phase is unobservable if $\phi = 0$. The effective weak phase may change due to non factorizable effects but the essential point here is that the weak phase ϕ can produce CP odd effects in nonleptonic B decays while its effect is hidden in B_s mixing. For the case of relative positive sign in $J_{sb,qq}$ the amplitude $\sim \sin \phi_c \cos \phi$. Note that if a weak phase is only associated with flavour changing $s - b$ vertex then $\phi_c = 0$ and there are no CP odd effects.

We can now look at the decay $B_d^0 \rightarrow \phi K_s$. For this decay we have to fierz the operators in Eq. 22 for the S/P model and we can define the Fierzed operators,

$$\begin{aligned}
H_{LL,S}^F &= -\frac{1}{2N_c} \bar{q}(1 - \gamma_5)b\bar{s}(1 - \gamma_5)q - \frac{1}{8N_c} \bar{q}\sigma_{\mu\nu}(1 - \gamma_5)b\bar{s}\sigma^{\mu\nu}(1 - \gamma_5)q, \\
H_{LR,S}^F &= -\frac{1}{2N_c} \bar{q}\gamma_\mu(1 - \gamma_5)b\bar{s}\gamma^\mu(1 + \gamma_5)q, \\
H_{RL,S}^F &= -\frac{1}{2N_c} \bar{q}\gamma_\mu(1 + \gamma_5)b\bar{s}\gamma^\mu(1 - \gamma_5)q, \\
H_{RR,S}^F &= -\frac{1}{2N_c} \bar{q}(1 + \gamma_5)b\bar{s}(1 + \gamma_5)q - \frac{1}{8N_c} \bar{q}\sigma_{\mu\nu}(1 + \gamma_5)b\bar{s}\sigma^{\mu\nu}(1 + \gamma_5)q,
\end{aligned} \tag{24}$$

where $q = s$ and we have done also a color Fierz and dropped octet operators that do not contribute in factorization.

We can therefore write,

$$\begin{aligned}
A(B_d^0 \rightarrow \phi K_s) &= A^{SM} + A_{\pm LR \pm RL}^{NP} + A_{LL+RR}^{NP}, \\
A^{SM} &= -\frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* Z \left[a_3^t + a_4^t + a_5^t - \frac{1}{2} a_7^t - \frac{1}{2} a_9^t - \frac{1}{2} a_{10}^t \right. \\
&\quad \left. - a_3^c - a_4^c - a_5^c + \frac{1}{2} a_7^c + \frac{1}{2} a_9^c + \frac{1}{2} a_{10}^c \right], \\
A_{\pm LR \pm RL}^{NP} &\sim \pm \langle K_s | \bar{s} \gamma_\mu b | B \rangle \langle \phi | \bar{s} \gamma^\mu s | 0 \rangle [e^{i\phi_-} + e^{i\phi_-}], \\
&\sim \pm \langle K_s | \bar{s} \gamma_\mu b | B \rangle \langle \phi | \bar{s} \gamma^\mu s | 0 \rangle \cos \phi_-, \\
A_{LL+RR}^{NP} &\sim \langle K_s | \bar{s} \sigma_{\mu\nu} b | B \rangle \langle \phi | \bar{s} \sigma^{\mu\nu} s | 0 \rangle [e^{i\phi_+} + e^{i\phi_+}], \\
&\sim \langle K_s | \bar{s} \sigma_{\mu\nu} b | B \rangle \langle \phi | \bar{s} \sigma^{\mu\nu} s | 0 \rangle \cos \phi_+, \\
Z &= 2f_\phi m_\phi F_{BK}(m_\phi^2) \varepsilon^* \cdot p_B,
\end{aligned} \tag{25}$$

where the SM contribution can be found in Ref. [26]. We therefore see that there are no new CP phases in $A(B_d \rightarrow \phi K_s)$. For the V/A models also it is easy to check that there are no NP phase in $B_d^0 \rightarrow \phi K_s$. This follows simply from the fact that only vector currents contribute to this decay and hence the matrix element of all the operators in $H_{eff}^{\Delta B=1}$ are equal. Explicitly, consider the decay $B_d^0 \rightarrow \phi K_s$ and using the currents associated with the relative negative sign in $J_{sb,qq}$, we can write

$$\begin{aligned}
\langle \phi K_s | H_{eff} | B_d^0 \rangle &\sim |\alpha| |c| \langle K_s | e^{i\phi} \bar{s} \gamma_\mu b - e^{-i\phi} \bar{s} \gamma_\mu b | B_d^0 \rangle \\
&\quad \langle \phi | e^{i\phi_c} \bar{s} \gamma^\mu s - e^{-i\phi_c} \bar{s} \gamma^\mu s | 0 \rangle, \\
&\sim 4|\alpha| |c| e^{i\pi/2} \sin \phi e^{i\pi/2} \sin \phi_c \langle K^- | \bar{s} \gamma_A b | B^- \rangle \\
&\quad \langle \pi^0 | \bar{d} \gamma^A \gamma_5 d | 0 \rangle, \\
&\sim 4|\alpha| |c| (-1) \sin \phi \sin \phi_c \langle K^- | \bar{s} \gamma_A b | B^- \rangle \\
&\quad \langle \pi^0 | \bar{d} \gamma^A \gamma_5 d | 0 \rangle.
\end{aligned} \tag{26}$$

This demonstrates that there are no new CP phases in $A(B_d^0 \rightarrow \phi K_s)$.

We now consider the case where $c_L \neq c_R$ in Eq. 21. For the case when $c_{L,R}$ are real there is a new weak phase of $\pi/2$ in $B^- \rightarrow K^- \pi^0$ for both the V/A and the S/P models. For the decay $B_d^0 \rightarrow \phi K_s$ a new weak phase of $\pi/2$ arises in V/A models while for the S/P models the phase in this decay depends on the size of $c_{L,R}$. Moving on to the most general case with complex $c_{L,R}$ the new weak phase in $B^- \rightarrow K^- \pi^0$ and $B_d^0 \rightarrow \phi K_s$ will depend on the phase in the $b \rightarrow s$ vertex as well as on the phases in $c_{L,R}$.

In the next section we will study a two higgs doublet model as an example of a NP model with $s - b$ interchange flavour symmetry.

3 A Specific NP Model

In Ref [22] we presented a two Higgs doublet model which has a 2-3 interchange flavour symmetry in the down quark sector like the $\mu - \tau$ interchange symmetry in the leptonic sector. The 2-3 symmetry is assumed in the gauge basis where the mass matrix has off diagonal terms and is fully 2-3 symmetric. Diagonalizing the mass matrix splits the masses of s and b or μ and τ and leads to vanishing $m_s(m_\mu)$. The breaking of the 2-3 symmetry is then introduced through the strange quark mass in the quark sector and the muon mass in the leptonic sector. The breaking of the 2-3 symmetry leads to flavor changing neutral currents (FCNC) in the quark sector and the charged lepton sector that are suppressed by $\frac{m_s}{m_b}$ and $\frac{m_\mu}{m_\tau}$ in addition to the mass of the Higgs boson of the second Higgs doublet. Additional FCNC effects of similar size can be generated from the breaking of the $s - b$ symmetry in the Yukawa coupling of the second Higgs doublet. A full discussion of the model is given in Ref [22] and we will repeat some of the discussions here for completeness.

We consider a Lagrangian of the form,

$$\mathcal{L}_Y^Q = Y_{ij}^U \bar{Q}_{i,L} \tilde{\phi}_1 U_{j,R} + Y_{ij}^D \bar{Q}_{i,L} \phi_1 D_{j,R} + S_{ij}^U \bar{Q}_{i,L} \tilde{\phi}_2 U_{j,R} + S_{ij}^D \bar{Q}_{i,L} \phi_2 D_{j,R} + h.c., \quad (27)$$

where ϕ_i , for $i = 1, 2$, are the two scalar doublets of a 2HDM, while $Y_{ij}^{U,D}$ and $S_{ij}^{U,D}$ are the non-diagonal matrices of the Yukawa couplings. After diagonalizing the Y matrix one can have FCNC couplings associated with the S matrix.

For convenience we express ϕ_1 and ϕ_2 in a suitable basis such that only the $Y_{ij}^{U,D}$ couplings generate the fermion masses. In such a basis one can write,

$$\langle \phi_1 \rangle = \begin{pmatrix} 0 \\ v/\sqrt{2} \end{pmatrix}, \quad \langle \phi_2 \rangle = 0. \quad (28)$$

The two Higgs doublets in this case are of the form,

$$\begin{aligned} \phi_1 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H^0 \end{pmatrix} + \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}\chi^+ \\ i\chi^0 \end{pmatrix}, \\ \phi_2 &= \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}H^+ \\ H^1 + iH^2 \end{pmatrix}. \end{aligned} \quad (29)$$

In principle there can be mixing among the neutral Higgs but here we neglect such mixing. We assume the doublet ϕ_1 corresponds to the scalar doublet of the SM and H^0 to the SM Higgs field. In addition, we assume that the second Higgs doublet does not couple to the up-type quarks ($S^U \equiv 0$). For the down type couplings in Eq. 27 we have,

$$\mathcal{L}_Y^D = Y_{ij}^D \bar{Q}_{i,L} \phi_1 D_{j,R} + S_{ij}^D \bar{Q}_{i,L} \phi_2 D_{j,R} + h.c. \quad (30)$$

We assume the following symmetries for the matrices Y^D and S^D :

- There is a discrete symmetry under which $d_{L,R} \rightarrow -d_{L,R}$
- There is a $s - b$ interchange symmetry: $s \leftrightarrow b$

The discrete symmetry involving the down quark is enforced to prevent $s \rightarrow d$ transition because of constraints from the kaon system. It also prevents $b \rightarrow d$ transitions since B_d mixing as well as the value of $\sin 2\beta$ measured in $B_d^0(t) \rightarrow J/\psi K_s$ are consistent with SM predictions. Although there may still be room for NP in $b \rightarrow d$ transitions, almost all deviations from the SM have been reported only in $b \rightarrow s$ transitions and so we assume no NP in $b \rightarrow d$ transitions in this work.

The above symmetries then give the following structure for the Yukawa matrices,

$$\begin{aligned} Y^D &= \begin{pmatrix} y_{11} & 0 & 0 \\ 0 & y_{22} & y_{23} \\ 0 & y_{23} & y_{22} \end{pmatrix}, \\ S^D &= \begin{pmatrix} s_{11} & 0 & 0 \\ 0 & s_{22} & s_{23} \\ 0 & s_{23} & s_{22} \end{pmatrix}. \end{aligned} \quad (31)$$

The down quark mass matrix, M^D is now given by $M^D = \frac{v}{\sqrt{2}}Y^D$. The matrix Y^D is symmetric and choosing the elements in Y^D to be real the mass matrix is diagonalized by,

$$\begin{aligned} M_{diag}^D &= U^T M^D U = \begin{pmatrix} \frac{v}{\sqrt{2}}y_{11} & 0 & 0 \\ 0 & \frac{v}{\sqrt{2}}(y_{22} - y_{23}) & 0 \\ 0 & 0 & \frac{v}{\sqrt{2}}(y_{22} + y_{23}) \end{pmatrix}, \\ U &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \end{aligned} \quad (32)$$

It is clear that the matrix U will also diagonalize the S^D matrix when we transform the quarks from the gauge to the mass eigenstate via $d_{L,R} \rightarrow U d_{L,R}$. Hence there are no FCNC effects involving the Higgs ϕ_2 .

The down quark masses are given by,

$$\begin{aligned} m_d &= \pm \frac{v}{\sqrt{2}}y_{11}, \\ m_s &= \pm \frac{v}{\sqrt{2}}(y_{22} - y_{23}), \\ m_b &= \pm \frac{v}{\sqrt{2}}(y_{22} + y_{23}). \end{aligned} \quad (33)$$

Since $m_s \ll m_b$ there has to be a fine tuned cancellation between y_{22} and y_{23} to produce the strange quark mass. Hence, it is more natural to consider the symmetry

limit $y_{22} = y_{23}$ which leads to $m_s = 0$. We then introduce the strange quark mass as a small breaking of the $s - b$ symmetry and consider the structure,

$$Y_n^D = \begin{pmatrix} y_{11} & 0 & 0 \\ 0 & y_{22}(1+2z) & y_{22} \\ 0 & y_{22} & y_{22} \end{pmatrix}, \quad (34)$$

with $z \sim 2m_s/m_b$ being a small number. Note that we do not break the $s - b$ symmetry in the 2 - 3 element so that the Y_n^D matrix remains symmetric. This down quark matrix is now diagonalized by,

$$M_{diag}^D = W^T M^D W = \begin{pmatrix} \pm \frac{v}{\sqrt{2}} y_{11} & 0 & 0 \\ 0 & \pm \frac{v}{\sqrt{2}} z y_{22} & 0 \\ 0 & 0 & \pm \frac{v}{\sqrt{2}} (2+z) y_{22} \end{pmatrix},$$

$$W = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{\sqrt{2}}(1 - \frac{1}{2}z) & \frac{1}{\sqrt{2}}(1 + \frac{1}{2}z) \\ 0 & \frac{1}{\sqrt{2}}(1 + \frac{1}{2}z) & \frac{1}{\sqrt{2}}(1 - \frac{1}{2}z) \end{pmatrix}. \quad (35)$$

In Eq. 35 we have dropped terms of $O(z^2)$. The down quark masses are now taken to be,

$$\begin{aligned} m_d &= \pm \frac{v}{\sqrt{2}} y_{11}, \\ m_s &= \pm \frac{v}{\sqrt{2}} z y_{22}, \\ m_b &= \pm \frac{v}{\sqrt{2}} (2+z) y_{22}. \end{aligned} \quad (36)$$

We find $z \approx \pm 2m_s/m_b$ and now the transformation to the mass eigenstate will generate FCNC effects involving ϕ_2 which is proportional to $z \sim 2m_s/m_b \sim \lambda^2$ where λ is the cosine of the Cabibbo angle. For definiteness we will choose the positive sign for z though both signs are allowed. At this point we can consider two scenarios: the first corresponds to the situation where the matrix S^D is still $s - b$ symmetric. This will result in $H_{NP,S}^{\Delta B=2}$ with $s - b$ symmetry and hence no observable phase in B_s mixing. In the second case we will allow for small breaking of the $s - b$ symmetry in S^D in Eq. 31. This will then result in $H_{NP,S}^{\Delta B=2}$ with broken $s - b$ symmetry and hence an observable phase in B_s mixing.

For the first case, S^D in the mass eigenstate basis has the form,

$$S^D \rightarrow S^{D'} = \begin{pmatrix} s_d e^{i\phi_{dd}} & 0 & 0 \\ 0 & s_s e^{i\phi_{ss}} & z s e^{i\phi_{sb}} \\ 0 & z s e^{i\phi_{sb}} & s_b e^{i\phi_{bb}} \end{pmatrix}, \quad (37)$$

where

$$s_d e^{i\phi_{dd}} = s_{11},$$

$$\begin{aligned}
s_s e^{i\phi_{ss}} &= (s_{22} - s_{23}), \\
s_b e^{i\phi_{bb}} &= (s_{22} + s_{23}), \\
s e^{i\phi_{sb}} &= s_{23}.
\end{aligned} \tag{38}$$

It is clear that $S^{D'}$ above, is symmetric under $s - b$ interchange.

The Hamiltonian describing the interaction of the Higgs, $H^{1,2}$, is given by,

$$\begin{aligned}
H_S &= \frac{1}{\sqrt{2}} [S_{sb}\bar{s}(1 + \gamma_5)b + S_{bs}\bar{b}(1 + \gamma_5)s] H^1 + h.c., \\
H_P &= i\frac{1}{\sqrt{2}} [S_{sb}\bar{s}(1 + \gamma_5)b + S_{bs}\bar{b}(1 + \gamma_5)s] H^2 + h.c.,
\end{aligned} \tag{39}$$

The relevant currents, J_{ij} are,

$$\begin{aligned}
J_{sb}^{S(P)} &= \frac{1}{\sqrt{2}} [S_{sb}\bar{s}(1 + \gamma_5)b \pm S_{bs}^*\bar{s}(1 - \gamma_5)b], \\
J_{qq}^{S(P)} &= \frac{1}{\sqrt{2}} [S_{qq}\bar{q}(1 + \gamma_5)q \pm S_{qq}^*\bar{q}(1 - \gamma_5)q],
\end{aligned} \tag{40}$$

where the $\pm$ sign correspond to scalar or pseudoscalar interactions. Using Eq. (37) we have,

$$S_{sb(bs)} = s e^{i\phi_{sb}} z, \tag{41}$$

$$S_{qq} = s_q e^{i\phi_{qq}}, \tag{42}$$

where $q = d, s$. After integrating out the heavy Higgs bosons, $H^{1,2}$, which we shall henceforth rename S and P bosons, we can generate the following effective Hamiltonian for $\Delta B = 2$ and $\Delta B = 1$ processes,

$$H_{NP}^{\Delta B=2} = \frac{1}{2m_{S(P)}^2} z^2 s^2 [\pm e^{i2\phi_{sb}} O_{RR} \pm e^{-i2\phi_{sb}} O_{LL} + O_{RL} + O_{LR}], \tag{43}$$

where O_{ab} are defined in Eq. 8 with $\gamma_A = 1$ and

$$\begin{aligned}
H_{eff}^{\Delta B=1} &= 4\frac{G_F}{\sqrt{2}} z \frac{s s_q}{g^2} \frac{m_W^2}{m_{S(P)}^2} [\pm e^{i\phi_+} H_{RR} \pm e^{-i\phi_+} H_{LL} \\
&\quad + e^{i\phi_-} H_{RL} + e^{-i\phi_-} H_{LR}],
\end{aligned} \tag{44}$$

where $\phi_{\pm} = \phi_{sb} \pm \phi_{qq}$, H_{ab} are defined in Eq. 22 with $\gamma_A = 1$ and the $\pm$ sign is for S and P exchange interactions. Note that $H_{NP}^{\Delta B=2}$ has the same structure as in Eq. 20 and is symmetric under $s - b$ interchange.

The expression for $M_{12}^{s,NP}$, obtained using the vacuum insertion approximation, is given as

$$\begin{aligned} M_{12}^{s,NP} &= \frac{1}{2m_{S(P)}^2} z^2 s^2 \left[(4 \mp \cos 2\phi_{sb} \frac{10}{3}) A + \frac{2}{3} \right] f_{B_s}^2 M_{B_s}, \\ A &= \frac{M_{B_s}^2}{(m_b + m_s)^2}. \end{aligned} \quad (45)$$

Here we assume that either the S or P interactions dominate and further we allow for a relative phase factor of ± 1 between $M_{12}^{s,NP}$ and $M_{12}^{s,SM}$. Following Ref. [7] we will take the SM prediction for ΔM_s^{SM} to be $\Delta M_s^{SM} = 23.4 ps^{-1}$ which then leads to $|1+\eta| = 0.74$ (see Eq. 5) allowing for values of $|\eta_{min}| = 0.26$ and $|\eta_{max}| = 1.74$. These values can be converted to values for $s_{min} \approx 0.1$ and $s_{max} \approx 0.25$ from Eq. 45 where we have chosen $\phi_{sb} = \pi/2$ for the pseudoscalar exchange interactions. This choice of ϕ_{sb} leads to the largest value for s . For our calculations we choose $m_s = 100$ MeV, $m_b = 4.8$ GeV, $f_{B_s} = 250$ MeV and $m_H = 1$ TeV. The choice for the higgs mass, m_H , is based on the fact that one expects new physics around $\sim$ TeV to stabilize the standard model higgs mass. We assume that the same new physics needed for the higgs mass stabilization is also responsible for new FCNC effects in B decays.

Let us now calculate the NP contribution to $B^- \rightarrow K^- \pi^0$. In $B \rightarrow \pi K$ decays, there are four classes of NP operators, differing in their color structure: $\bar{s}_\alpha \Gamma_i b_\alpha \bar{q}_\beta \Gamma_j q_\beta$ and $\bar{s}_\alpha \Gamma_i b_\beta \bar{q}_\beta \Gamma_j q_\alpha$ ($q = u, d$). The matrix elements of these operators are then combined into single NP amplitudes, denoted by $\mathcal{A}'^q e^{i\Phi'_q}$ and $\mathcal{A}'^{C,q} e^{i\Phi_q^C}$ respectively with $q = u, d$ [14, 27]. We have

$$\begin{aligned} \langle K^- \pi^0 | H_{eff}^{\Delta B=1} | B^- \rangle &= |\mathcal{A}'^{comb}| e^{i\Phi'} = 4 \frac{G_F}{\sqrt{2}} \chi_s A_{dd} [\pm 2i \sin \phi_+ - 2i \sin \phi_-], \\ &= 16 \frac{G_F}{\sqrt{2}} \chi_s A_{dd} e^{i\pi/2} \begin{cases} \cos \phi_{sb} \sin \phi_{dd} & H = S, \\ -\sin \phi_{sb} \cos \phi_{dd} & H = P, \end{cases} \\ \chi_s &= \frac{2m_s}{m_b} \frac{ss_d}{g^2} \frac{m_W^2}{m_{S(P)}^2}, \\ A_{dd} &= \langle K^- | \bar{s}b | B^- \rangle \langle \pi^0 | \bar{d}\gamma_5 d | 0 \rangle. \end{aligned} \quad (46)$$

Here $\mathcal{A}'^{comb} e^{i\Phi'} \equiv -\mathcal{A}'^u e^{i\Phi'_u} + \mathcal{A}'^d e^{i\Phi'_d}$. In the case of scalar exchange the CP odd effect in the amplitude is due to the phase ϕ_{dd} while in the case of the pseudoscalar exchange the CP odd effect comes from the phase ϕ_{sb} which also appears in B_s mixing. Hence we will consider the pseudoscalar exchange only for our quantitative analysis.

We first note that we can choose, in Eq. 46, the phase $\Phi' = 90^\circ$ which is then consistent with a fit to the $B \rightarrow K\pi$ data found in Ref. [14] which found $\Phi' \sim 100^\circ$ and $|\mathcal{A}'^{comb}/T'| = 1.64$ where T' is the SM tree amplitude. Here we will attempt to see if this value for the ratio $|\mathcal{A}'^{comb}/T'|$ is consistent with the ΔM_s measurements.

For the pseudoscalar exchange model we can write for the ratio $|\mathcal{A}'^{comb}/T'|$, using naive factorization[21],

$$|\mathcal{A}'^{comb}/T'| = \left| \frac{16 \sin \phi_{sb} \cos \phi_{dd} \chi_s A_{dd}}{(c_1 + c_2/N_C) V_{ub}^* V_{us} \langle \pi^0 | \bar{b} \gamma_\mu (1 - \gamma_5) u | B^- \rangle \langle K^- | \bar{u} \gamma^\mu (1 - \gamma_5) s | 0 \rangle} \right|, \quad (47)$$

where $c_{1,2}$ are the Wilson's coefficients of the SM effective Hamiltonian. We can rewrite the above expression as,

$$|\mathcal{A}'^{comb}/T'| = |16 \sin \phi_{sb} \cos \phi_{dd} \chi_s| M, \quad M = \left| \frac{[(m_B^2 - m_K^2)/(m_b - m_s)] F_0^K (m_\pi^2/2m_d) f_\pi/\sqrt{2}}{f_K(m_B^2 - m_\pi^2) F_0^\pi/\sqrt{2} \left(c_1 + \frac{c_2}{N_C}\right) |V_{ub}^* V_{us}|} \right| \quad (48)$$

We take $(f_K/f_\pi)(F_0^\pi/F_0^K) \sim 1$, $|V_{ub}^* V_{us}/V_{tb}^* V_{ts}| = 1/48$ and $c_1 + c_2/N_C = 1.018$. Now using the values of s obtained from B_s mixing we obtain the maximum possible values $|\mathcal{A}'^{comb}/T'| \approx (0.47, 1.2)s_d(6\text{MeV}/m_d)$. Assuming $s_d \sim 1$ in Eq. 46, we find the lesser value for $|\mathcal{A}'^{comb}/T'|$ a bit smaller than the fit value in Ref. [14] of 1.64. One could increase $|\mathcal{A}'^{comb}/T'|$ by choosing s_d large enough, $s_d \sim 3$, which would indicate a strongly coupled theory. The second, larger value for $|\mathcal{A}'^{comb}/T'|$ is in better agreement with the fit in Ref. [14]. One could also increase $|\mathcal{A}'^{comb}/T'|$ by lowering m_d which can be taken to be in the range 4-8 MeV [28]. Hence the predictions for the size of NP in $B \rightarrow K\pi$ decays does not rule out the scenario that there is no new phase in B_s mixing. However a measurement of a new weak phase in $B_d \rightarrow \phi K_s$ would indicate new phase in B_s mixing in this model.

To introduce a phase in B_s mixing we have to break the $s-b$ interchange flavour symmetry. This can be done by allowing S^D in Eq. 31 to break the $s-b$ interchange symmetry. We choose,

$$S^D = \begin{pmatrix} s_{11} & 0 & 0 \\ 0 & s_{22} & s_{23}(1+2\epsilon) \\ 0 & s_{23} & s_{22}(1+2\delta) \end{pmatrix}, \quad (49)$$

with ϵ, δ being small quantities of the same size or smaller than z . The transformation to the mass eigenstate leads to the general parametrization,

$$S^D \rightarrow S^{D'} = \begin{pmatrix} s_d e^{i\phi_{dd}} & 0 & 0 \\ 0 & s_s e^{i\phi_{ss}} & z s e^{i\phi_{sb}} - \epsilon p e^{i\psi_{sb}} \\ 0 & z s e^{i\phi_{sb}} + \epsilon p e^{i\psi_{sb}} & s_b e^{i\phi_{bb}} \end{pmatrix}, \quad (50)$$

where

$$\begin{aligned}
s_d e^{i\phi_{dd}} &= s_{11}, \\
s_s e^{i\phi_{ss}} &= (s_{22} - s_{23} + s_{22}\delta) + s_{22}\delta z - s_{23}\epsilon, \\
s_b e^{i\phi_{bb}} &= (s_{22} + s_{23} + s_{22}\delta) - s_{22}\delta z + s_{23}\epsilon, \\
s e^{i\phi_{sb}} &= s_{22} \frac{\delta}{z} + s_{23}, \\
p e^{i\psi_{sb}} &= s_{23}.
\end{aligned} \tag{51}$$

There is now an additional FCNC involving $b \rightarrow s$ transitions whose source is the $s-b$ symmetry breaking in S^D . Note that the 2–3 off-diagonal elements in $S^{D'}$ contain a part that is symmetric under $s-b$ interchange and a part that is antisymmetric under the $s-b$ interchange. The parameters in $S^{D'}$ can be obtained or constrained from a fit to B decay data. A phase in B_s mixing arises from the interference of the the $s-b$ symmetric and the $s-b$ antisymmetric piece.

The structure in Eq. 50 is rather complicated and for the illustration of how a new phase in B_s mixing may arise we can reduce the parameters in the model to simplify the discussion. Hence, we will assume $\delta = 0$ which leads to $\phi_{sb} = \psi_{sb}$, $s = p$ and calling $s_{23} \equiv s$ we obtain,

$$S^D \rightarrow S^{D'} = \begin{pmatrix} s_d e^{i\phi_{dd}} & 0 & 0 \\ 0 & s_s e^{i\phi_{ss}} & s(z - \epsilon) e^{i\phi_{sb}} \\ 0 & s(z + \epsilon) e^{i\phi_{sb}} & s_b e^{i\phi_{bb}} \end{pmatrix}, \tag{52}$$

The relevant currents, J_{ij} are,

$$\begin{aligned}
J_{sb}^{S(P)} &= \frac{1}{\sqrt{2}} s \left[(z - \epsilon) e^{i\phi_{sb}} \bar{s} (1 + \gamma_5) b \pm (z + \epsilon) e^{-i\phi_{sb}} \bar{s} (1 - \gamma_5) b \right], \\
J_{qq}^{S(P)} &= \frac{1}{\sqrt{2}} \left[S_{qq} \bar{q} (1 + \gamma_5) q \pm S_{qq}^* \bar{q} (1 - \gamma_5) q \right],
\end{aligned} \tag{53}$$

where the $\pm$ sign correspond to scalar or pseudoscalar interactions. The B_s mixing is now generated by,

$$H_{NP}^{\Delta B=2} = \frac{1}{2m_{S(P)}^2} z^2 s^2 \left[\pm (1 - \frac{\epsilon}{z})^2 e^{i2\phi_{sb}} O_{RR} \pm (1 + \frac{\epsilon}{z})^2 e^{-i2\phi_{sb}} O_{LL} + (1 - \frac{\epsilon^2}{z^2}) (O_{RL} + O_{LR}) \right]. \tag{54}$$

Now we can try two approximations. One with $\frac{\epsilon}{z}$ small and one with $\epsilon \sim z$. The possibility $\epsilon \gg z$ is ruled out by the $B \rightarrow K\pi$ data as it does not produce a new weak phase [22]. For $\frac{\epsilon}{z}$ small, we get keeping only terms linear in $\frac{\epsilon}{z}$,

$$H_{NP}^{\Delta B=2} =$$

$$\frac{1}{2m_{S(P)}^2} z^2 s^2 \left[\pm e^{i2\phi_{sb}} O_{RR} \pm e^{-i2\phi_{sb}} O_{LL} \pm \frac{2\epsilon}{z} (e^{-i2\phi_{sb}} O_{LL} - e^{+i2\phi_{sb}} O_{RR}) + (O_{RL} + O_{LR}) \right]. \quad (55)$$

We can now write $M_{12}^{s,NP}$ as,

$$\begin{aligned} M_{12}^{s,NP} &= \frac{2}{m_{S(P)}^2} z^2 s^2 \left[\left(1 \mp \frac{5}{6} \cos 2\phi_{sb}\right) A + \frac{2}{12} \pm i \frac{5}{3} \frac{\epsilon}{z} \sin 2\phi_{sb} A \right] f_{B_s}^2 M_{B_s}, \\ A &= \frac{M_{B_s}^2}{(m_b + m_s)^2}. \end{aligned} \quad (56)$$

So we see that there is an observable phase in B_s mixing which is proportional to $\frac{\epsilon}{z}$ and is given by,

$$\begin{aligned} \sin \phi_{B_s} &= \frac{\text{Im} M_{12}^{s,NP}}{|M_{12}^{s,NP}|}, \\ &\approx \frac{\frac{5}{3} \frac{\epsilon}{z} \sin 2\phi_{sb} A}{\left(1 \mp \frac{5}{6} \cos 2\phi_{sb}\right) A + \frac{2}{12}}. \end{aligned} \quad (57)$$

We now study the predictions for $b \rightarrow s\bar{q}q$ transitions. To simplify the discussion we will choose S_{qq} to be real and so any new phase in $b \rightarrow s\bar{q}q$ is due to the weak phase in J_{sb} . It is also clear that only the pseudoscalar exchange contribute to $B^- \rightarrow K^- \pi^0$. One can check that there will be a new weak phase in $B_d^0 \rightarrow \phi K_s$ proportional to $\frac{\epsilon}{z}$ while the weak phase in $B^- \rightarrow K^- \pi^0$ will deviate from $\pi/2$ by an amount proportional to $\frac{\epsilon}{z}$.

For $\epsilon = z$, The relevant currents, J_{ij} are now given by,

$$\begin{aligned} J_{sb}^{S(P)} &= \pm \frac{1}{\sqrt{2}} s z e^{-i\phi_{sb}} \bar{s} (1 - \gamma_5) b, \\ J_{qq}^{S(P)} &= \frac{1}{\sqrt{2}} s_q [\bar{q} (1 + \gamma_5) q \pm \bar{q} (1 - \gamma_5) q], \end{aligned} \quad (58)$$

where again we have chosen $S_{qq} = s_q$ to be real. It is now clear that $H_{NP}^{\Delta B=2} \sim J_{sb}^2$ has an observable phase which can be large and large NP phase can appear in $B_d \rightarrow \phi K_s$ and $B^- \rightarrow K^- \pi^0$. It is easy to check that the phase in $B^- \rightarrow K^- \pi^0$ and $B_d^0 \rightarrow \phi K_s$ is half the phase in B_s mixing and hence if the B_s mixing is observed to be small it would indicate a small new weak phase in $B^- \rightarrow K^- \pi^0$ and in $B_d^0 \rightarrow \phi K_s$ in contrast to the case considered above. In passing we note that if $\phi_{sb} = \pi$ then a NP phase of $\pi/2$ can appear in $B^- \rightarrow K^- \pi^0$ but a phase will also appear in the decay $B_d^0 \rightarrow \phi K_s$ in contrast to the $s - b$ symmetric case considered above.

To summarize, if the B_s mixing phase is observed to be small and is due to the small breaking of the $s - b$ interchange symmetry then one should observe a large new CP odd phase, close to $\pi/2$, in the decay $B^- \rightarrow K^- \pi^0$ and a small new CP odd

phase in $B_d^0 \rightarrow \phi K_s$. On the other hand if the B_s mixing is small due to small NP phase in the $b \rightarrow s$ current then we should observe a small NP phase in the decay $B^- \rightarrow K^- \pi^0$ and in $B_d^0 \rightarrow \phi K_s$.

4 Conclusions

In summary, in this paper we have studied the implication of the ΔM_s measurement on $b \rightarrow s\bar{q}q$ transitions. We pointed out that in a fairly general class of NP the new physics phase of B_s mixing may be unobservable if there is a $s - b$ interchange flavour symmetry even in the presence of new CP odd phase in the $b \rightarrow s$ transition. This phase can however appear in certain $b \rightarrow s\bar{q}q$ transitions like $B \rightarrow K\pi$ decays but not in others like the $B_d^0 \rightarrow \phi K_s$ decay. Working within a two higgs doublet model we calculated the allowed NP contribution to $B \rightarrow K\pi$ decays with the new ΔM_s measurement. We also showed that if the NP weak phase is only in the $b \rightarrow s$ current and a small observed B_s mixing phase is due to the small breaking of the $s - b$ interchange symmetry then one should observe a large new CP odd phase, close to $\pi/2$, in the decay $B^- \rightarrow K^- \pi^0$ but a small CP odd phase in $B_d^0 \rightarrow \phi K_s$. On the other hand a small B_s mixing phase and a small NP phase in the decays $B^- \rightarrow K^- \pi^0$ and $B_d^0 \rightarrow \phi K_s$ would indicate the absence of the $s - b$ flavour symmetry.

Acknowledgements: This work is financially supported by NSERC of Canada. We thank R.N. Mohapatra for useful discussion.

References

- [1] V. Abazov [D0 Collaboration], arXiv:hep-ex/0603029.
- [2] G. Gomez-Ceballos [CDF Collaboration], talk at FPCP 2006, <http://fpcp2006.triumf.ca/talks/day3/1500/fpcp2006.pdf>
- [3] M. Bona *et al.* [UTfit Collaboration], JHEP **0507**, 028 (2005); updated results available at <http://utfit.roma1.infn.it/>.
- [4] J. Charles *et al.* [CKMfitter Group], Eur. Phys. J. C **41**, 1 (2005); updated results available at <http://ckmfitter.in2p3.fr/>.
- [5] M. Endo and S. Mishima, arXiv:hep-ph/0603251; M. Blanke, A. J. Buras, D. Guadagnoli and C. Tarantino, arXiv:hep-ph/0604057; J. Foster, K. i. Okumura and L. Roszkowski, arXiv:hep-ph/0604121; K. Kleinknecht and B. Renk, arXiv:hep-ph/0604202; K. Cheung, C. W. Chiang, N. G. Deshpande and J. Jiang, arXiv:hep-ph/0604223; G. Isidori and P. Paradisi, arXiv:hep-ph/0605012; S. Khalil, arXiv:hep-ph/0605021.

- [6] Z. Ligeti, M. Papucci and G. Perez, arXiv:hep-ph/0604112.
- [7] P. Ball and R. Fleischer, arXiv:hep-ph/0604249.
- [8] M. Beneke, G. Buchalla, A. Lenz and U. Nierste, Phys. Lett. B **576**, 173 (2003) [arXiv:hep-ph/0307344]; M. Ciuchini, E. Franco, V. Lubicz, F. Mescia and C. Tarantino, JHEP **0308**, 031 (2003) [arXiv:hep-ph/0308029].
- [9] Y. Grossman, Y. Nir and G. Raz, arXiv:hep-ph/0605028.
- [10] B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0507087; B. Aubert *et al.* [BABAR Collaboration], [hep-ex/0503011]; B. Aubert *et al.* [BABAR Collaboration], [hep-ex/0502019]; B. Aubert *et al.* [BABAR Collaboration], [hep-ex/0502017]; B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **94**, 161803 (2005) [arXiv:hep-ex/0408127].
- [11] K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0507037;
K. F. Chen *et al.* [Belle Collaboration], Phys. Rev. D **72**, 012004 (2005) [arXiv:hep-ex/0504023]; K. Abe *et al.* [BELLE Collaboration], arXiv:hep-ex/0409049.
- [12] E. Barberio *et al.* [HFAG], arXiv:hep-ex/0603003; updated results available at <http://www.slac.stanford.edu/xorg/hfag/>.
- [13] Experiment: B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **89**, 281802 (2002); arXiv:hep-ex/0408062, arXiv:hep-ex/0408080, arXiv:hep-ex/0408081; A. Bornheim *et al.* [CLEO Collaboration], Phys. Rev. D **68**, 052002 (2003); Y. Chao *et al.* [Belle Collaboration], Phys. Rev. D **69**, 111102 (2004). Theory: A. J. Buras, R. Fleischer, S. Recksiegel and F. Schwab, Eur. Phys. J. C **32**, 45 (2003), Phys. Rev. Lett. **92**, 101804 (2004), Nucl. Phys. B **697**, 133 (2004), arXiv:hep-ph/0410407; V. Barger, C. W. Chiang, P. Langacker and H. S. Lee, Phys. Lett. B **598**, 218 (2004); S. Mishima and T. Yoshikawa, arXiv:hep-ph/0408090; Y. L. Wu and Y. F. Zhou, arXiv:hep-ph/0409221; H. Y. Cheng, C. K. Chua and A. Soni, arXiv:hep-ph/0409317; Y. Y. Charng and H. n. Li, arXiv:hep-ph/0410005; X. G. He and B. H. J. McKellar, arXiv:hep-ph/0410098; C. S. Kim, S. Oh and C. Yu, arXiv:hep-ph/0505060.
- [14] S. Baek, P. Hamel, D. London, A. Datta and D. A. Suprun, Phys. Rev. D **71**, 057502 (2005) [arXiv:hep-ph/0412086].
- [15] T. E. Browder, A. Datta, X. G. He and S. Pakvasa, Phys. Rev. D **57**, 6829 (1998) [arXiv:hep-ph/9705320]; A. Datta, X. G. He and S. Pakvasa, Phys. Lett. B **419**, 369 (1998) [arXiv:hep-ph/9707259].
- [16] For a study of triple products in the SM and with new physics, see A. Datta and D. London, Int. J. Mod. Phys. A **19**, 2505 (2004).

- [17] B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0408017. Note that the earlier Belle measurements of the same quantities do not show any signs of a nonzero triple-product asymmetry, see K.-F. Chen *et al.* [Belle Collaboration], Phys. Rev. Lett. **91**, 201801 (2003).
- [18] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **91**, 171802 (2003); K. F. Chen *et al.* [BELLE Collaboration], arXiv:hep-ex/0503013.
- [19] Particle Data Group, Ref. [28]; B. Aubert [BABAR Collaboration], arXiv:hep-ex/0408093; K. Senyo [Belle Collaboration].
- [20] J. Smith, talk presented at FPCP 2006, <http://fpcp2006.triumf.ca/talks/day4/1030/fpcp.pdf>
- [21] S. Baek, A. Datta, P. Hamel, O. F. Hernandez and D. London, arXiv:hep-ph/0508149.
- [22] A. Datta and P. J. O'Donnell, Phys. Rev. D **72**, 113002 (2005) [arXiv:hep-ph/0508314].
- [23] T. Inami and C.S. Lim, Progr. Theor. Phys. **65** (1981) 297.
- [24] C. S. Lam, hep-ph/0104116; T. Kitabayashi and M. Yasue, Phys.Rev. **D67** 015006 (2003); W. Grimus and L. Lavoura, hep-ph/0305046; 0309050; Y. Koide, Phys.Rev. **D69**, 093001 (2004); for examples of such theories, see W. Grimus and L. Lavoura, hep-ph/0305046; 0309050.
- [25] R. N. Mohapatra, SLAC Summer Inst. lecture; <http://www-conf.slac.stanford.edu/ssi/2004>; hep-ph/0408187; JHEP, **0410**, 027 (2004); W. Grimus, A. S.Joshipura, S. Kaneko, L. Lavoura, H. Sawanaka, M. Tanimoto, hep-ph/0408123; A. Ghosal, Mod. Phys. Lett. A **19**, 2579 (2004).
- [26] A. Datta, Phys. Rev. D **66**, 071702 (2002) [arXiv:hep-ph/0208016].
- [27] A. Datta, M. Imbeault, D. London, V. Page, N. Sinha and R. Sinha, Phys. Rev. D **71**, 096002 (2005); A. Datta and D. London, Phys. Lett. B **595**, 453 (2004) [arXiv:hep-ph/0404130].
- [28] S. Eidelman *et al.* [Particle Data Group Collaboration], Phys. Lett. B **592** (2004) 1, <http://pdg.lbl.gov/pdg.html>.